

Assessing capacity for diagnosing tuberculosis in children in sub-Saharan African HIV care settings

M. J. A. Reid,*† S. Saito,† R. Fayorsey,† R. J. Carter,† E. J. Abrams†

* Botswana University of Pennsylvania Partnership, University of Pennsylvania, Philadelphia, Pennsylvania, † Mailman School of Public Health, International Center for AIDS Care and Treatment Programs, Columbia University, New York, New York, USA

SUMMARY

Research on the prevalence of pediatric-specific tuberculosis (TB) diagnostics in sub-Saharan Africa is scarce. We assessed the availability of pediatric TB diagnostic tests at 651 pediatric human immunodeficiency virus care and treatment sites across nine African countries: 54% of the sites had access to sputum culture capacity and 51% to chest X-ray services. While 87% of sites

had access to smear microscopy, only 6% had the capacity to perform sputum induction and 5% to perform gastric aspirate. These findings confirm that diagnostic resources for the accurate diagnosis of pediatric TB are limited. Capacity-building initiatives to improve sputum collection in children are urgently required.

KEY WORDS: tuberculosis; HIV; pediatric; diagnosis

THE DIAGNOSIS of tuberculosis (TB) in children living with the human immunodeficiency virus (HIV) in sub-Saharan Africa is challenging;¹ sputum and culture confirmation rarely exceeds 30–40%.² In most instances, the diagnosis of TB is based on clinical criteria alone.³ The clinical difficulties of diagnosing TB in HIV-positive children are well described. However, accurate diagnosis is also limited by programmatic limitations: specifically, the limited capability to acquire sputum samples in children unable to expectorate and the lack of trained personnel to perform and interpret TB diagnostic procedures in children.

In the present analysis, we describe the availability of TB diagnostic procedures and tests at sites providing pediatric HIV services in nine sub-Saharan countries. We sought to determine the proportion of sites that had access to chest X-ray (CXR), sputum culture and smear microscopy, and how many reported a capacity to perform sputum induction, naso-pharyngeal aspirate (NPA) and gastric aspirate (GA). We also sought to determine facility characteristics associated with the presence of these diagnostic capabilities.

METHODS

In September 2010, we surveyed 651 sites providing HIV services in nine countries (Cote d'Ivoire, Ethiopia, Kenya, Mozambique, Nigeria, Rwanda, South Africa, Swaziland and Tanzania) that received sup-

port from the International Center for AIDS Care and Treatment Programs (ICAP) at Columbia University, New York, through funding from the government of the United States. National policy in each of these countries includes routine screening for TB in HIV-infected children.

Survey questions included access to TB diagnostic tests (smear microscopy, CXR and sputum culture onsite or offsite), capacity to perform specific pediatric TB diagnostic procedures (sputum induction, GA and NPA) and program characteristics. All sites provided HIV services to HIV-infected and HIV-exposed children aged 0–15 years. A recent assessment of this annual survey showed 83% test-retest agreement across multiple countries.⁴ χ^2 tests were used to assess the association of program and facility characteristics with the capacity to perform or with access to the following pediatric TB diagnostic procedures or tests: smear microscopy, GA, sputum induction, NPA, sputum culture and CXR. We examined the presence of these procedures or tests in relation to location (urban, semi-urban, rural, defined as per Demographic and Health Surveys criteria⁵), facility type (primary, secondary, tertiary and private), program maturity (years since initiation of pediatric HIV services), and overall program size (cumulative number of children and adults enrolled in care). Statistical analysis was performed using SAS software version 9.2 (SAS, Cary, NC, USA).

Correspondence to: Michael J A Reid, HIV Division, Botswana UPenn Partnership, PO Box AC 157 ACH, Gaborone, Botswana. Tel: (+267) 724 787 77. Fax: (+267) 317 0957. e-mail: michael.reid@doctors.org.uk; reidm@botswana-upenn.co.bw

Article submitted 8 December 2011. Final version accepted 1 February 2012.

[A version in French of this article is available from the Editorial Office in Paris and from the Union website www.theunion.org]

The protocol was reviewed by the Institutional Review Board of the Columbia University, New York, and received non-human subject research determination.

RESULTS

As of September 2010, 651 sites surveyed in this analysis had cumulatively enrolled 82 413 HIV-infected and HIV-exposed children (Table 1). Sputum smear microscopy was available at 87% of the sites, serving 98% of all children enrolled in care ($n = 565$ sites, range across countries [RAC] 28–100%); 5% ($n = 34$, RAC 0–21%) had the capacity to perform GA collection, 6% ($n = 39$, RAC 0–24%) had the capacity to perform sputum induction and 2% ($n = 12$, RAC 0–7%) had the capacity to perform NPA. Mycobacterial culture (performed onsite or offsite) was available at 54% ($n = 352$, RAC 2–97%) of the sites. Nearly all sites with advanced capacity to ob-

tain sputum (sputum induction and GA) performed sputum culture; however, respectively only 10% and 8% of the sites with access to sputum culture had sputum induction or GA capacity; 51% ($n = 330$, RAC 18–89%) had access to CXR, performed either onsite or offsite.

In bivariate analysis, availability of TB diagnostics (Table 2) and the capacity to procure sputum samples (Table 3) were more likely at urban than rural sites, with the exception of sputum induction. Tertiary and secondary sites were more likely to have the capacity to perform GA (4% vs. 10%, $P = 0.0037$) and sputum induction (3% vs. 13%, $P < 0.001$) and have access to CXR (36% vs. 84%, $P < 0.001$) compared to primary care sites. CXR (33% vs. 69%, $P < 0.001$) was more prevalent at sites with larger HIV programs.

DISCUSSION

The findings of our analysis are cause for alarm, given the high toll of disease and death among HIV-infected children in Africa. While the majority of the sites had the capacity to perform smear microscopy and had access to mycobacterial culture, essential for diagnosing TB in *both* adults and children, a much smaller fraction had capacity to perform any of the sputum collection procedures necessary in children unable to expectorate.

Given the limitations of clinical criteria alone in diagnosing TB in children with HIV,^{6,7} it is worrying that relatively few sites have the capacity to obtain pediatric sputum samples for microscopy and culture. Lack of resources was particularly acute at primary care sites and at sites located in rural areas. These findings highlight the urgent need to increase access to accurate TB diagnostics in areas of high HIV prevalence and TB endemicity. To be effective, scale-up in access needs to occur commensurate with the policy of decentralizing HIV services from tertiary/secondary to primary care sites that is ongoing in many sub-Saharan countries.

There are several exciting developments in the field of TB diagnostics, including Cepheid's GeneXpert system, Xpert[®] MTB/RIF (Sunnyvale, CA, USA).⁸ However, the accuracy of this assay in diagnosing TB in children is predicated on obtaining good quality sputum samples. The utility of Xpert MTB/RIF will be compromised if sputum collection capacity is not improved. Our analysis demonstrates that any expansion in Xpert MTB/RIF use must occur in tandem with concerted efforts to build capacity for improved sputum collection for those children unable to expectorate spontaneously.

This study had several limitations. We were not able to distinguish whether lack of diagnostic capacity was due to lack of human resources or diagnostic hardware. Furthermore, where it was available, we

Table 1 HIV care and treatment site characteristics, September 2010 ($N = 651$)

	Number of pediatric HIV care and treatment sites n (%)	Children aged 0–14 years enrolled in HIV care and treatment* n (%)
Total	651	82 413
Country		
Cote d'Ivoire	60 (9)	455 (1)
Ethiopia	62 (10)	8 145 (10)
Kenya	157 (24)	15 510 (19)
Mozambique	60 (9)	23 138 (28)
Nigeria	28 (4)	2 863 (3)
Rwanda	42 (6)	3 742 (4)
South Africa	67 (10)	8 666 (11)
Swaziland	49 (8)	14 883 (18)
Tanzania	126 (19)	5 011 (6)
Location		
Urban	128 (20)	30 987 (38)
Semi-urban	204 (31)	28 101 (34)
Rural	319 (49)	23 327 (28)
Clinic type		
Primary	379 (58)	27 673 (34)
Secondary/tertiary	196 (30)	50 769 (62)
Private/other	76 (12)	3 971 (4)
Program size (cumulative number of HIV-infected adults and children enrolled in care)		
<352	320 (50)	4 926 (6)
≥ 352	324 (50)	77 487 (94)
Data missing	7	0
Program maturity (years since inception of pediatric HIV services), years		
≥ 5	50 (8)	22 735 (28)
≥ 3 –<5	155 (24)	27 616 (34)
≥ 1 –<3	358 (55)	30 855 (37)
<1	88 (14)	1 207 (1)

*Cumulative number enrolled, pre-ART and on ART. HIV = human immunodeficiency virus; ART = antiretroviral therapy.

Table 2 Bivariate analysis: factors associated with availability of diagnostic tests for diagnosing TB in children at pediatric HIV care and treatment sites in nine countries, September 2010 (N = 651)*

	Smear microscopy			Sputum culture			Chest X-ray		
	n (%)	PR (95%CI)	P value	n (%)	PR (95%CI)	P value	n (%)	PR (95%CI)	P value
Total pediatric HIV sites	565 (87)			352 (54)			330 (51)		
Total pediatric patients served by surveyed sites	80 596 (98)			61 471 (7)			60 967 (74)		
Location									
Urban	107 (92)	1.1 (0.99–1.1)	NS	78 (67)	1.2 (1.0–1.4)	NS	89 (77)	2.2 (1.8–2.6)	<0.0001 [†]
Semi-urban	174 (83)	0.96 (0.89–1.0)	NS	87 (42)	0.73 (0.60–0.87)	NS	126 (60)	1.7 (1.4–2.1)	<0.0001 [†]
Rural	284 (87)	1.0		187 (57)	1.0		115 (35)	1.0	
Facility type									
Primary	317 (83)	1.0		206 (54)	1.0		137 (36)	1.0	
Secondary/tertiary	189 (98)	1.2 (1.1–1.2)	<0.0001 [†]	126 (66)	1.2 (1.1–1.4)	0.0059 [†]	161 (84)	2.3 (2.0–2.7)	<0.0001 [†]
Private/other	59 (76)	0.91 (0.80–1.0)	NS	20 (26)	0.47 (0.32–0.70)	0.0002	32 (41)	1.1 (0.85–1.5)	NS
Program size (cumulative no. of children and/or adults enrolled)									
<352	244 (76)	0.97 (0.96–0.98)	<0.0001 [†]	121 (38)	0.95 (0.93–0.97)	<0.0001 [†]	105 (33)	1.0	<0.0001 [†]
≥352	316 (98)	1.0		230 (71)	1.0		224 (69)	2.1 (1.8–2.5)	
Program maturity (years since initiation of pediatric HIV services), years									
≥5	49 (98)	1.3 (1.1–1.5)	0.0004	39 (78)	1.9 (1.4–2.6)	<0.0001	44 (88)	2.8 (2.0–3.8)	<0.0001 [†]
≥3–<5	155 (100)	1.0 (0–0)	NS	110 (71)	1.7 (1.3–2.3)	<0.0001 [†]	113 (73)	2.3 (1.7–3.2)	<0.0001 [†]
≥1–<3	299 (84)	1.1 (1.0–1.3)	0.0452 [†]	167 (47)	1.1 (0.87–1.50)	NS	145 (41)	1.3 (0.91–1.8)	NS
<1	62 (70)	1.0		36 (41)	1.0		28 (32)	1.0	
Country									
Cote d'Ivoire	17 (3)			1 (2)			19 (32)		
Ethiopia	62 (11)			37 (60)			55 (89)		
Kenya	147 (26)			135 (86)			58 (37)		
Mozambique	57 (10)			52 (87)			49 (82)		
Nigeria	28 (5)			10 (36)			24 (86)		
Rwanda	42 (7)			31 (74)			15 (36)		
South Africa	66 (12)			65 (97)			43 (64)		
Swaziland	37 (7)			10 (20)			9 (18)		
Tanzania	109 (19)			11 (9)			58 (46)		

*The table excludes missing values. Categories are not mutually exclusive. Facilities were able to choose multiple responses.

[†]Statistically significant.

TB = tuberculosis; HIV = human immunodeficiency virus; PR = prevalence ratio; CI = confidence interval; NS = non-significant.

were unable to differentiate the quality, comprehensiveness or availability of diagnostic capacity. Strengths of our study include the large number of sites, which allowed us to conduct an analysis of the availability of pediatric TB diagnostics across several sub-Saharan countries. Given that ICAP provides antiretroviral drugs (ARVs) to approximately 9% of persons on ARVs in eight of the nine countries included,* we believe that our findings are arguably representative of President's Emergency Plan for AIDS Relief (PEPFAR) funded programs from a diverse array of settings and contexts.⁹

* Swaziland excluded.

CONCLUSION

Across 651 pediatric HIV sites in Africa, resources for the diagnosis of TB in children were limited. While 87% of the sites had access to smear microscopy, fewer than 10% had the capacity to procure sputum samples in children unable to expectorate spontaneously. Capacity-building initiatives to improve sputum collection in children are urgently needed.

Acknowledgements

The authors thank and acknowledge the field staff from the International Center for AIDS Care and Treatment Programs who completed the site assessments. They also acknowledge the Ministries of Health of Kenya, Tanzania, Ethiopia, Rwanda, Mozambique, South Africa, Nigeria, Lesotho, Democratic Republic of Congo, Zambia,

Table 3 Bivariate analysis: factors associated with reported capacity to procure sputum samples in children at pediatric HIV care and treatment sites in nine countries, September 2010 (*n* = 651)*

	Naso-pharyngeal aspirate			Gastric aspirates			Induced sputum		
	<i>n</i> (%)	PR (95%CI)	<i>P</i> value	<i>n</i> (%)	PR (95%CI)	<i>P</i> value	<i>n</i> (%)	PR (95%CI)	<i>P</i> value
Total pediatric HIV sites	12 (2)			34 (5)			39 (6)		
Total pediatric patients served by surveyed sites	3 814 (5)			11 236 (14)			7 729 (9)		
Location									
Urban	6 (5)	16.9 (2.1–138.6)	0.0086 [†]	14 (12)	3.9 (1.8–8.6)	0.0006 [†]	7 (6)	1.0 (0.43–2.3)	NS
Semi-urban	5 (2)	7.8 (2.1–66.3)	0.0599 [†]	10 (5)	1.6 (0.66–3.7)	NS	12 (6)	0.94 (0.47–1.9)	NS
Rural	1 (0.3)	1.0		10 (3)	1.0		20 (6)	1.0	
Facility type									
Primary	5 (1)	1.0		14 (4)	1.0		12 (3)	1.0	
Secondary/tertiary	7 (4)	2.8 (0.89–8.6)	NS	19 (10)	2.7 (1.4–5.3)	0.0037 [†]	24 (13)	4.0 (2.0–7.8)	<0.0001
Private/other	0	—	—	1 (1)	0.35 (0.05–2.6)	NS	3 (4)	1.2 (0.35–4.2)	NS
Program size (cumulative no. of children and/or adults enrolled)									
<352	2 (1)	4.9 (1.1–22.4)	0.038 [†]	8 (3)	1.0		14 (4)	1.0	
≥352	10 (3)	1.0		26 (8)	3.2 (1.5–7.0)	0.0033 [†]	25 (8)	1.8 (0.93–3.3)	NS
Program maturity (years since initiation of pediatric HIV services), years									
≥5	8 (16)			9 (18)	5.3 (1.5–18.6)	0.0096	8 (16)	1.8 (0.70–4.4)	NS
≥3–<5	3 (2)			10 (6)	1.9 (0.53–3.7)	NS	10 (6)	0.71 (0.29–1.7)	NS
≥1–<3	1 (0)			12 (3)	1.0 (0.28–3.4)	NS	13 (4)	0.40 (0.17–0.93)	0.0342 [†]
<1	0			3 (3)	1.0		8 (9)	1.0	
Country									
Cote d'Ivoire	1 (2)			0			0		
Ethiopia	0			0			15 (24)		
Kenya	0			2 (1)			11 (7)		
Mozambique	3 (5)			4 (7)			2 (3)		
Nigeria	1 (4)			3 (11)			1 (4)		
Rwanda	3 (7)			8 (19)			3 (7)		
South Africa	3 (4)			14 (21)			3 (4)		
Swaziland	0			2 (4)			2 (4)		
Tanzania	1 (1)			1 (1)			2 (2)		

*The table excludes missing values. Categories are not mutually exclusive. Facilities were able to choose multiple responses.

[†]Statistically significant.

TB = tuberculosis; HIV = human immunodeficiency virus; PR = prevalence ratio; CI = confidence interval; NS = non-significant.

Swaziland and Cote d'Ivoire, whose HIV care and treatment programs they endeavor to support through this work. Finally, the authors thank W El Sadr for overall leadership on program activities. The activities described in this article were funded through the President's Emergency Plan for AIDS Relief through the US Centers for Disease Control and Prevention.

References

- Hesseling A C, Schaaf H S, Gie R P, Starke J R, Beyers N. A critical review of diagnostic approaches used in the diagnosis of childhood tuberculosis. *Int J Tuberc Lung Dis* 2002; 6: 1038–1045.
- Zar H J, Hanslo D, Apolles P, Swingler G, Hussey G. Induced sputum versus gastric lavage for microbiological confirmation of pulmonary tuberculosis in infants and young children: a prospective study. *Lancet* 2005; 365: 130–134.
- Harries A D, Parry C, Nyong'onya Mbewe L, et al. The pattern of tuberculosis in Queen Elizabeth Central Hospital, Blantyre, Malawi: 1986–1995. *Int J Tuberc Lung Dis* 1997; 1: 346–351.
- International Center for AIDS Care and Treatment Programs. Unpublished PFACTs data quality assurance exercise from nine African countries. New York, NY, USA: ICAP, 2010.
- United States Agency for International Development. Demographic and health surveys, 2012. Calverton, MD, USA: DHS, 2012. <http://www.measuredhs.com/> Accessed March 2012.
- Walters E, Cotton M F, Rabie H, Schaaf H S, Walters L O, Marais B J. Clinical presentation and outcome of tuberculosis in human immunodeficiency virus infected children on anti-retroviral therapy. *BMC Pediatr* 2008; 8: 1.
- Marais B J, Graham S M, Cotton M F, Beyers N. Diagnostic and management challenges for childhood tuberculosis in the era of HIV. *J Infect Dis* 2007; 196 (Suppl 1): S76–S85.
- Nicol M P, Workman L, Isaacs W, et al. Accuracy of the Xpert MTB/RIF test for the diagnosis of pulmonary tuberculosis in children admitted to hospital in Cape Town, South Africa: a descriptive study. *Lancet Infect Dis* 2011; 11: 819–824.
- World Health Organization/Joint United Nations Programme on HIV/AIDS/United Nations International Children's Emergency Fund. Towards universal access: scaling up priority HIV/AIDS interventions in the health sector. Geneva, Switzerland: WHO, 2010. <http://www.who.int/hiv/pub/2010progressreport/report/en/index.html> Accessed March 2012.

R É S U M É

En Afrique sub-saharienne, il n'y a que peu de recherche concernant la prévalence des techniques de diagnostic de la tuberculose (TB) spécifiques à la pédiatrie. Nous avons évalué la disponibilité de tests de diagnostic TB pédiatriques dans 651 sites de soins et de traitement du virus de l'immunodéficience humaine pédiatrique dans neuf pays d'Afrique. La possibilité de culture des crachats existait dans 54% des sites et des services de radio-

graphie thoracique dans 51% des sites. Alors que l'examen microscopique des frottis est disponible dans 87% des sites, 6% seulement sont aptes à réaliser l'induction des crachats et 5% le tubage gastrique. Ces observations confirment que les ressources diagnostiques sont limitées pour le diagnostic précis de la TB pédiatrique. Des initiatives de formation à une amélioration de la collecte des crachats chez les enfants sont nécessaires d'urgence.

R E S U M E N

Son muy escasas las investigaciones que describen la utilización de métodos diagnósticos de la tuberculosis (TB) que sean específicos de los niños en África subsahariana. En el presente estudio se evaluó la disponibilidad de pruebas diagnósticas de la TB en niños en 651 centros pediátricos de atención y tratamiento de la infección por el virus de la inmunodeficiencia humana en nueve países africanos. El 54% de los centros contaba con acceso al cultivo del esputo y 51% a los servicios de radiografía

de tórax. El 87% de las instituciones podía ofrecer la baciloscopia del esputo, pero en solo 6% se podía practicar la inducción del esputo y en 5% el aspirado gástrico. Estos resultados confirman la insuficiencia de los recursos diagnósticos existentes que permitan establecer un diagnóstico preciso de TB en los niños. Se precisan con urgencia iniciativas de fortalecimiento de las capacidades, encaminadas a mejorar la calidad de la recogida de muestras de esputo en los niños.
